



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\sum_{n=1}^{\infty} \frac{U_n}{r^n} = \frac{1}{\sqrt{5}} \left( \frac{a + b\rho_1}{\rho_1(r - \rho_1)} - \frac{1}{\sqrt{5}} \frac{a + b\rho_2}{\rho_2(r - \rho_2)} \right).$$

The right-hand member now reduces to

$$\frac{b + ar - a}{r^2 - r - 1}$$

when  $\rho_1$  is replaced by  $(1 + \sqrt{5})/2$  and  $\rho_2$  by  $(1 - \sqrt{5})/2$ .

*Note.*—We are publishing this solution for the reason that the previously published solution referred to did not consider the question of convergency correctly, and the proper investigation of this question was the Proposer's chief reason for proposing the problem. EDITORS.

#### ALGEBRA.

##### 429. Proposed by C. N. SCHMALL, New York City.

It is given that  $d_1, d_2, d_3$  are the greatest common divisors of  $y$  and  $z$ ,  $z$  and  $x$ ,  $x$  and  $y$ , respectively; also that  $m_1, m_2, m_3$  are the least common multiples of the same pairs of members. If  $d$  and  $m$  are the greatest common divisor and least common multiple, respectively, of  $x, y$ , and  $z$ , show that

$$\frac{m}{d} = \left( \frac{m_1 m_2 m_3}{d_1 d_2 d_3} \right)^{\frac{1}{2}}.$$

SOLUTION BY FRANK IRWIN, University of California.

It is evident that we can get the least common multiple of two numbers by dividing their product by their greatest common divisor:

$$m_1 = \frac{yz}{d_1}, \quad m_2 = \frac{xz}{d_2}, \quad m_3 = \frac{xy}{d_3}.$$

Similarly with the three numbers  $x, y, z$ , if we divide their product by  $d_1 d_2 d_3$ , we should have their least common multiple, except that we have divided out  $d$  once too often:

$$m = \frac{xyz}{d_1 d_2 d_3} \cdot d.$$

We have then:

$$\left( \frac{m_1}{d_1} \cdot \frac{m_2}{d_2} \cdot \frac{m_3}{d_3} \right)^{\frac{1}{2}} = \left( \frac{yz}{d_1^2} \cdot \frac{zx}{d_2^2} \cdot \frac{xy}{d_3^2} \right)^{\frac{1}{2}} = \frac{xyz}{d_1 d_2 d_3} = \frac{m}{d}.$$

Also solved by A. H. HOLMES, ELMER SCHUYLER, G. W. HARTWELL, FRANK R. MORRIS, N. P. PANDYA, HERBERT N. CARLETON, PAUL CAPRON, J. A. CAPARO, and the PROPOSER.

#### GEOMETRY.

##### 443. Proposed by C. N. SCHMALL, New York City.

A quadrilateral of any shape whatever is divided by a transversal into two quadrilaterals. The diagonals of the original figure and those of the two resulting (smaller) figures are then drawn. Show that their three points of intersection are collinear.

#### III. SOLUTION BY LAENAS G. WELD, Pullman, Ills.

To the triangle  $ABC$  draw the transversals  $MN$ , intersecting  $AB$  in  $M$  and  $AC$  in  $N$ , and  $QR$  intersecting  $AB$  in  $Q$  and  $AC$  in  $R$ . Then  $BCRQ$ ,  $BCNM$  and